

A NEW FAMILY OF POLYNOMIAL IDENTITIES FOR COMPUTING DETERMINANTS

GEORGY P. EGORYCHEV

ABSTRACT. We give new definitions for the determinant over commutative ring K , noncommutative ring $\mathbf{K}$, noncommutative ring $\mathcal{K}$ with associative powers, over noncommutative nonassociative ring $\mathfrak{K}$, and study their properties.

Let K be a commutative ring, $\mathbf{K}$ a noncommutative associative ring, $\mathcal{K}$ a noncommutative ring with associative powers (one-monomial associativity), and $\mathfrak{K}$ be a noncommutative nonassociative ring; let each ring be with division by integers. Here we obtain a new family of polynomial identities for determinant over the ring K , which allows us to give new definitions for determinants over rings $\mathbf{K}$, $\mathcal{K}$, $\mathfrak{K}$, and to study their properties. These definitions are closely related to the definition of symmetrized *Barvinok's* determinant $s\det(A)$ [2] different from the well-known determinant of Dieudonné over a division ring, the quasideterminant [6], and other well-known determinants over noncommutative associative rings [1]. We also estimate the computational complexity of the obtained formulas for the determinants.

Let $A = (a_{ij})$ be an $n \times n$ matrix with elements from the ring K . Let S_n be the set of all permutations $\sigma = (\sigma(1), \dots, \sigma(n))$ of the set $\{1, \dots, n\}$, $\tau(\sigma)$ be the number of inversions in σ . Let $S_n^{(e)}$ and $S_n^{(o)}$ be the subsets of even and odd permutations in S_n , respectively. We call the sequence of elements of $a_{1\sigma(1)}, \dots, a_{n\sigma(n)}$ the diagonal $l(\sigma)$ of matrix A , the sequence of elements $a_{i_1\sigma(i_1)}, \dots, a_{i_k\sigma(i_k)}$, $1 \leq i_1 < \dots < i_k \leq n$ the subdiagonal l of length k of matrix A . We denote by $L_k^{(e)} \left(L_k^{(o)} \right)$ the set of all subdiagonals of length k of diagonals from $S_n^{(e)} \left(S_n^{(o)} \right)$, $k = 1, \dots, n$. The function $su(l)$ on the subdiagonal (diagonal) $l = \text{diag}(a_{i_1\sigma(i_1)}, \dots, a_{i_k\sigma(i_k)})$, is defined to be the sum of its elements, i.e. $su(l) := \sum_{s=1}^k a_{i_s\sigma(i_s)}$.

Definition 1. (*the determinant over a commutative ring*)

$$\det(A) := \sum_{\sigma \in S_n} (-1)^{\tau(\sigma)} a_{1\sigma(1)} \dots a_{n\sigma(n)}.$$

In the following Main Theorem we employ the well-known polarization theorem [3, 4] to obtain a new family of polynomial identities for determinants that contain up to $n!$ free variables.

Date: December 3, 2011.

2010 Mathematics Subject Classification. Primary 11C20, 16Rxx; Secondary 15A15.

Key words and phrases. determinant, theory of rings, polynomial identities.

Theorem 1 (Main Theorem). *If $A = (a_{ij})$ is an $n \times n$ matrix over the ring K , then the following formulas hold:*

(0.1)

$$\det(A) = \frac{1}{n!} \sum_{\sigma \in S_n} (-1)^{\tau(\sigma)} \left\{ (-1)^n \gamma_\sigma^n + \sum_{k=1}^n ((-1)^{n-k} \sum_{1 \leq j_1 < \dots, j_k \leq n} (\gamma_\sigma + \sum_{s=1}^k a_{j_s \sigma(j_s)})^n) \right\},$$

where $\{\gamma_\sigma\}_{\sigma \in S_n}$ is the set of $n!$ free variables $\gamma_\sigma \in K$. In particular for all $\gamma_\sigma = \gamma$ and for all $\gamma_\sigma = 0$ we have, correspondingly

$$\begin{aligned} \det(A) &= \frac{(-1)^{n-1}}{n!} \left\{ \sum_{l \in L_{n-1}^{(e)}} (\gamma + su(l))^n - \sum_{l \in L_{n-1}^{(o)}} (\gamma + su(l))^n + \right. \\ &\quad \left. + \sum_{l \in L_n^{(o)}} (\gamma + su(l))^n - \sum_{l \in L_n^{(e)}} (\gamma + su(l))^n \right\}, \end{aligned} \quad (0.2)$$

(0.3)

$$\det(A) = \frac{(-1)^{n-1}}{n!} \left\{ \sum_{l \in L_{n-1}^{(e)}} su^n(l) - \sum_{l \in L_{n-1}^{(o)}} su^n(l) + \sum_{l \in L_n^{(o)}} su^n(l) - \sum_{l \in L_n^{(e)}} su^n(l) \right\}.$$

Corollary 1. *If $A = (a_{ij})$ is an $n \times n$ matrix over $\mathbb{C}$ then for $t = 1, 2, \dots, n-1$ the following identities are hold:*

$$(0.4) \quad \sum_{l \in L_{n-1}^{(e)}} su^t(l) - \sum_{l \in L_{n-1}^{(o)}} su^t(l) + \sum_{l \in L_n^{(o)}} su^t(l) - \sum_{l \in L_n^{(e)}} su^t(l) = 0.$$

Corollary 2. *(a new criterion for independence of matrix rows (columns)). If $A = (a_{ij})$ is an $n \times n$ matrix over the ring K , then $\det(A) = 0$ iff the following identity holds*

$$\sum_{l \in L_n^{(e)}} su^n(l) - \sum_{l \in L_n^{(o)}} su^n(l) = \sum_{l \in L_{n-1}^{(e)}} su^n(l) - \sum_{l \in L_{n-1}^{(o)}} su^n(l).$$

Remark 1. a) The formulas (0.1) – (0.3) are obtained using all characteristic properties of the determinant [9].

b) Formula (0.1) for $n!$ free variables $\{\gamma_\sigma\}_{\sigma \in S_n}$ generates $2^{n!}$ different polynomial identities if we put each $\gamma_\sigma = 0$ or $\gamma_\sigma \neq 0$. Each of these formulas can be taken as the definition for $\det(A)$, it requires its own number of arithmetical operations when calculated and according to Corollary 1 generates a new set of identities.

c) Formula (0.1) and its special cases employ (besides divisions by $n!$) only operations $+$, $-$, and raising to power of n , yet does not use a commutativity of multiplication in the ring K .

Definition 2. *(the noncommutative determinant $\text{edet}(A)$ over ring $\mathbf{K}$ or $\mathcal{K}$).*

(0.5)

$$\text{edet}(A) := \frac{(-1)^{n-1}}{n!} \left\{ \sum_{l \in L_{n-1}^{(e)}} su^n(l) - \sum_{l \in L_{n-1}^{(o)}} su^n(l) + \sum_{l \in L_n^{(o)}} su^n(l) - \sum_{l \in L_n^{(e)}} su^n(l) \right\}.$$

Lemma 1. *(the properties of $\text{edet}(A)$ over ring $\mathbf{K}$).*

a) $\text{edet}(A)$ is a polyadditive function of the rows and columns of matrix A ;

b) $\text{edet}(A)$ is an antisymmetric function of the rows and columns of matrix A ;

- c) $\text{edet}(A) = 0$, if the matrix A contains zero row or zero column;
d) $\text{edet}(A) = 0$, if the matrix A contains two equal rows (columns);
e) $\text{edet}(A) = \text{edet}(A^T)$.

The Laplace formulas and the formula $\det(AB) = \det(A) \times \det(B)$ are not valid in the general case.

Let $\text{sdet}(A)$ be the symmetrized Barvinok's determinant over ring $\mathbf{K}$ [2]:

$$\text{sdet}(A) := \frac{1}{n!} \sum_{\mu \in S_n} \sum_{\sigma \in S_n} (-1)^{\tau(\sigma) + \tau(\mu)} a_{\mu(1)\sigma(1)} \dots a_{\mu(n)\sigma(n)}.$$

Lemma 2. The following formulas are valid for $\text{edet}(A)$ over ring $\mathbf{K}$:

$$\text{sdet}(A) = \text{edet}(A),$$

$\text{edet}(I_n) = e$, if the ring $\mathbf{K}$ contains the unit e , and I_n is the identity $n \times n$ matrix.

Lemma 3. (computational complexity of $\text{edet}(A)$). The computation of $\text{edet}(A)$ via formulas (0.3) and (0.5) for large n requires of order $(3n-1) \times n!$ additions and $(n+1)! \times \ln n$ multiplications (see [8])

Formula (0.5) allows to introduce the following new general definition

Definition 3. (the determinant $\text{edet}(A)$ over ring $\mathfrak{K}$).

$$\text{edet}(A) := \frac{(-1)^{n-1}}{n!} \left\{ \sum_{l \in L_{n-1}^{(e)}} \text{Ass}(su^n(l)) - \sum_{l \in L_{n-1}^{(o)}} \text{Ass}(su^n(l)) + \sum_{l \in L_n^{(o)}} \text{Ass}(su^n(l)) - \sum_{l \in L_n^{(e)}} \text{Ass}(su^n(l)) \right\},$$

where $\text{Ass}(a^n)$, $a \in \mathfrak{K}$, is associate operator

$$\text{Ass}(a^n) := \{(a \times (a \times (a \dots (a \times a)))) + \dots + (((a \times a) \times a) \times a) \dots \times a\} / C_n,$$

and $C_n = (2n-2)!/n!(n-1)!$ is the Catalan number of n -products in nonassociative algebra (all arrangements of brackets in the n -product $a \times a \times \dots \times a$).

Lemma 4. (the properties of the determinant $\text{edet}(A)$ over nonassociative rings $\mathcal{K}$ and $\mathfrak{K}$). The determinant $\text{edet}(A)$ over rings $\mathcal{K}$ and $\mathfrak{K}$ satisfies properties b) – e) of Lemma 1. The Laplace formulas, the polyadditivity by rows and columns of matrix A and the formula $\text{edet}(AB) = \text{edet}(A) \times \text{edet}(B)$ are not valid in general case.

Obtaining similar results is of interest for Shur functions, the mixed discriminants and many other matrix functions of planar and space matrices related to determinants. Our results may find applications in the theory of permanents [4], the theory of n -Lie algebras [5], differential geometry [10] and elsewhere.

The author is grateful to his colleagues M.N. Davletshin, V.M. Kopytov, Y.N. Nuzhin, A.V. Timofeenko, I.P. Shestakov for discussion of the basic results of this work and a number of useful remarks.

REFERENCES

- [1] Arvind V. and Srinivasa S. On the Hardness of Noncommutative Determinant, Electronic Colloquium on Computational Complexity, Report No. 103, 2009.
- [2] Barvinok A. New permanent estimators via non-commutative determinants. Arxiv preprint math/0007153, 2000, arXiv:math/0007153.
- [3] Cartan H. Elementary theory of analytic functions of one or several complex variables. Dover Publ., New York, 1995, 228 p.

- [4] Egorychev G.P. Discrete mathematics. Permanents. Krasnoyarsk, Siberian Federal University, 2007, 272 p., (transl. in English: Springer, 2011).
- [5] Filippov V.T. On the n -Lie algebra of Jacobians, *Sibirsk. Mat. Zh.*, 39:3, 1998, 660–669.
- [6] Gel'fand I. M. and Retakh V. S. Determinants of matrices over noncommutative rings, *Funktsional. Anal. i Prilozhen.* 25, 1991, No. 2, 13–25, 96.
- [7] Zevlakov K.A., Slin'ko A.M., Shestakov I.P. and Shirshov A.I. Rings that are nearly associative, *Acad. Press*, 1982, 432 p.
- [8] Kochergin V.V. About complexity of computation one-terms of powers, *Discrete analysis*, IM SO RAN, v. 27, Novosibirsk, 1994, 94 – 107 (in Russian)
- [9] Muir T. A Treatise on the Theory of Determinants, New York: Dover, 1960.
- [10] Sabinin L.V. Methods of Nonassociative Algebra in Differential Geometry. in Supplement to Russian translation of S.K. Kobayashi and K. Nomizu "Foundations of Differential Geometry", 1, Moscow, Nauka, 1981.

RUSSIA, KRASNOYARSK,

E-mail address: `anott@scn.ru`